



# Open Research Online

---

The Open University's repository of research publications  
and other research outputs

## Reassessing the stable isotope composition assigned to methane flux from natural wetlands in isotope-constrained budgets

### Conference or Workshop Item

#### How to cite:

Hornibrook, Edward; Maxfield, Peter; Gauci, Vincent and Stott, Andrew (2013). Reassessing the stable isotope composition assigned to methane flux from natural wetlands in isotope-constrained budgets. In: European Geosciences Union General Assembly 2013, 7-12 Apr 2013, Vienna, Austria.

For guidance on citations see [FAQs](#).

© 2013 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<http://meetingorganizer.copernicus.org/EGU2013/EGU2013-13060.pdf>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)



## Reassessing the stable isotope composition assigned to methane flux from natural wetlands in isotope-constrained budgets

Edward Hornibrook (1), Peter Maxfield (2), Vincent Gauci (3), and Andrew Stott (4)

(1) University of Bristol, School of Earth Sciences, Bristol, UK (ed.hornibrook@bristol.ac.uk), (2) Department of Applied Sciences, University of the West of England, Frenchay Campus, Coldharbour Lane, Bristol, BS16 1QY, UK (pete.maxfield@uwe.ac.uk), (3) Centre for Earth, Planetary, Space and Astronomical Research (CEPSAR), The Open University, Milton Keynes, MK7 6AA, UK (v.gauci@open.ac.uk), (4) Centre for Ecology & Hydrology, CEH-Lancaster, Lancaster Environment Centre, Library Avenue Bailrigg, Lancaster, LA1 4AP UK (astott@ceh.ac.uk)

Stable isotope ratios in  $\text{CH}_4$  preserve information about its origin and history, and are commonly used to constrain global  $\text{CH}_4$  budgets. Wetlands are key contributors to the atmospheric burden of  $\text{CH}_4$  and typically are assigned a stable carbon isotope composition of  $\sim -60$  permil in isotope-weighted stable isotope models despite the considerable range of  $\delta^{13}\text{C}(\text{CH}_4)$  values ( $\sim -100$  to  $-40$  permil) known to occur in these diverse ecosystems. Kinetic isotope effects (KIEs) associated with the metabolism of  $\text{CH}_4$ -producing microorganisms generate much of the natural variation but highly negative and positive  $\delta^{13}\text{C}(\text{CH}_4)$  values generally result from secondary processes (e.g., diffusive transport or oxidation by soil methanotrophs). Despite these complexities, consistent patterns exist in the isotope composition of wetland  $\text{CH}_4$  that can be linked conclusively to trophic status and consequently, natural succession or human perturbations that impact nutrient levels.

Another challenge for accurate representation of wetlands in carbon cycle models is parameterisation of sporadic  $\text{CH}_4$  emission events. Abrupt release of large volumes of  $\text{CH}_4$ -rich bubbles in short periods of time can account for a significant proportion of the annual  $\text{CH}_4$  flux from a wetland but such events are difficult to detect using conventional methods. New infrared spectroscopy techniques capable of high temporal resolution measurements of  $\text{CH}_4$  concentration and stable isotope composition can readily quantify short-lived  $\text{CH}_4$  pulses. Moreover, the isotope data can be used conclusively to determine shifts in the mode of  $\text{CH}_4$  transport and provide the potential to link initiation of abrupt emission events to forcing by internal or external factors.